

Scaling limit for a drainage network model

C. F. Coletti^{*§} L. R. G. Fontes^{†§} E. S. Dias^{‡§}

Abstract We consider the two dimensional version of a drainage network model introduced by Gangopadhyay, Roy and Sarkar, and show that the appropriately rescaled family of its paths converges in distribution to the Brownian web. We do so by verifying the convergence criteria proposed by Fontes, Isopi, Newman and Ravishankar.

Keywords and phrases Drainage networks, coalescing random walks, Brownian web, coalescing Brownian motions

2000 Mathematics Subject Classification 60K35, 60K40, 60F17

1 Introduction and results

The Two-dimensional Drainage Network Model Let $\mathbb{Z}$ be the set of the integers and $\Omega = (\omega(z), z \in \mathbb{Z}^2)$ be a family of Bernoulli independent random variables with parameter $p \in (0, 1)$. Let $\mathbb{P}_p$ and $\mathbb{E}_p$ be the probability and expectation induced by those variables in the product space $(\{0, 1\})^{\mathbb{Z}^2}$. Consider a second family $\Upsilon = (v(z), z \in \mathbb{Z}^2)$ of Bernoulli independent random variables with parameter $1/2$. Let $\mathbb{P}_{1/2}$ and $\mathbb{E}_{1/2}$ be the probability and expectation induced by these variables in the product space $(\{0, 1\})^{\mathbb{Z}^2}$. Let $\mathbb{P} = \mathbb{P}_p \times \mathbb{P}_{1/2}$ be the product probability induced by the Bernoulli variables in the product space $\{0, 1\}^{\mathbb{Z}^2} \times \{0, 1\}^{\mathbb{Z}^2}$ and let $\mathbb{E}$ be the expectation operator with respect to this probability.

For $z = (z_1, z_2) \in \mathbb{Z}^2$, say that z' is in the next level if $z' = (w, z_2 + 1)$ for some $w \in \mathbb{Z}$. Let $h(z)$ be the closest open vertex to z in the next level with respect to the distance induced by the l_1 norm. That is to say, $\omega(h(z)) = 1$ and $\sum_{i=1}^2 |h(z)_i - z_i| = \min\{\sum_{i=1}^2 |x_i - z_i| : \omega(x) = 1 \text{ and } x_2 = z_2 + 1\}$. If there exist two closest open vertex to z in the next level, the connection will be to its left if $v(z) = 0$ and to its right if $v(z) = 1$. Now, let $h^0(z) = z$ and iteratively, for $n \geq 1$, $h^n(z) = h(h^{n-1}(z))$.

Now let $\mathcal{G} = (V, \mathcal{E})$ be the random directed graph with vertices $V = \mathbb{Z}^2$ and edges $\mathcal{E} = \{(u, h(u)) : u \in \mathbb{Z}^2\}$. This model was proposed in [1] and will be

^{*}Research supported by FAPESP grant 2006/54511-2

[†]Partially supported by CNPq grants 307978/2004-4 and 484351/2006-0, and FAPESP grant 2004/07276-2

[‡]Research supported by FAPESP grant 2004/13008-0

[§]Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010, Cidade Universitária, 05508-090 São Paulo SP, Brasil, emails: {cristian, lrenato, dias}@ime.usp.br

called the *Gangopadhyay, Roy and Sarkar* (two-dimensional drainage network) *model*, or GRS model, for short.

Main result It follows from the construction that the GRS model may be seen as a set of continuous paths, as follows. For any $z = (z_1, z_2) \in \mathbb{Z}^2$ we define the path $X^z = \{X^z(s), s \geq z_2\}$ in $\mathbb{R}^2$ as the linearly interpolated line composed by all edges $\{(h^k(z), h^{k+1}(z)) : k \in \mathbb{N}\}$ of the model, with $X^z(k) = h^k(z)$, $k \in \mathbb{N}$, where h^k is the k -th composition of h , h^0 meaning the identity. Clearly, X^z is a continuous path starting at time z_2 . We Let

$$\mathcal{X} := \{X^z : z \in \mathbb{Z}^2\}, \quad (1.1)$$

which we also call the *drainage network*, and consider its diffusive rescaling

$$\mathcal{X}_\delta := \{(\delta x_1, \delta^2 x_2) \in \mathbb{R}^2 : (x_1, x_2) \in \mathcal{X}\}, \quad (1.2)$$

for $\delta \in (0, 1]$. Our main result below shows that $\mathcal{X}_\delta$ converges in distribution to the Brownian web.

Several authors constructed random processes that formally correspond to coalescing one-dimensional Brownian motions starting from every space-time point [2, 3, 4, 5, 6]. In [5], the Brownian web is characterized as a random variable taking values in a metric space whose points are compact sets of paths. Denote by $(\mathcal{H}, d_{\mathcal{H}})$ the complete separable metric space where the Brownian web is defined. Denote also by $\mathcal{F}_{\mathcal{H}}$ the corresponding Borel σ -algebra generated by $d_{\mathcal{H}}$.

The closure in path space of the rescaled drainage network $\mathcal{X}_\delta$, also denoted by $\mathcal{X}_\delta$, is an $(\mathcal{H}, \mathcal{F}_{\mathcal{H}})$ -valued random variable.

Theorem 1 *The rescaled drainage network $\mathcal{X}_\delta$ converges in distribution to the Brownian web as $\delta \rightarrow 0$.*

In order to prove Theorem 1, we will verify the convergence criteria of [5]. To describe them, we need the following definition. Given $t_0 \in \mathbb{R}$, $t > 0$, $a < b$, and a $(\mathcal{H}, \mathcal{F}_{\mathcal{H}})$ -valued random variable $\mathcal{V}$, let $\eta_{\mathcal{V}}(t_0, t; a, b)$ be the $\{0, 1, 2, \dots, \infty\}$ -valued random variable giving the number of *distinct* points in $\mathbb{R} \times \{t_0 + t\}$ that are touched by paths in $\mathcal{V}$ which also touch some point in $[a, b] \times \{t_0\}$.

We can now state the convergence criteria. Let $\mathcal{D}$ be a countable dense set of points in $\mathbb{R}^2$.

Theorem 2 ([5]) *Suppose that $\mathcal{X}_1, \mathcal{X}_2, \dots$ are $(\mathcal{H}, \mathcal{F}_{\mathcal{H}})$ -valued random variables with noncrossing paths. If, in addition, the following three conditions are valid, the distribution of $\mathcal{X}_n$ converges to the distribution of the standard Brownian web.*

(I₁) *There exist $\theta_n^y \in \mathcal{X}_n$ such that for any deterministic $y_1, \dots, y_m \in \mathcal{D}$, $\theta_n^{y_1}, \dots, \theta_n^{y_m}$ converge in distribution as $n \rightarrow \infty$ to coalescing Brownian motions (with unit diffusion constant) starting at $y_1, \dots, y_m$.*

(B₁) $\limsup_{n \rightarrow \infty} \sup_{(a, t_0) \in \mathbb{R}^2} \mathbb{P}(\eta_{\mathcal{X}_n}(t_0, t; a, a + \epsilon) \geq 2) \rightarrow 0$ as $\epsilon \rightarrow 0+$;

(B₂) $\epsilon^{-1} \limsup_{n \rightarrow \infty} \sup_{(a, t_0) \in \mathbb{R}^2} \mathbb{P}(\eta_{\mathcal{X}_n}(t_0, t; a, a + \epsilon) \geq 3) \rightarrow 0$ as $\epsilon \rightarrow 0+$.

The present model can be viewed as a discrete space, long range version of the two-dimensional Poisson tree introduced in [7]. In [8], the weak convergence to the Brownian web of the rescaled collection of paths of that model was established. The approach is the same as the one pursued here. There are considerable technical differences in the present work, though. The main ones are as follows: a) the interaction among the paths in the present model is a long range one, differently from the former one; this requires considerably more care in the proofs of (I_1) and of an estimate on the tail of the distribution of the coalescence time of two given paths, which we use in the proof of (B_2) (see Theorems 8 and 4), and b) as in [8], we resort to an FKG type of argument in an estimation in the proof of (B_2) , and for that we need to establish a monotonicity property of certain conditional probabilities on conditioning paths (see Proposition 6); this is also more delicate here than in [8].

Another drainage network for which the rescaled paths were shown to converge in distribution to the Brownian web is the one proposed by Scheidegger [9], which is equivalent to coalescing simple random walks. This case was treated in [5]. In the general context of drainage networks, our results may be seen as part of the scaling theory for those models. See [10].

The GRS model may be seen also as coalescing random walks, in this case whose paths are non-crossing and have unbounded dependence among themselves. The other networks mentioned above may also be seen as non-crossing coalescing random walks. Coalescing non-simple random walks which are independent before coalescence, and whose paths may thus cross one another, were studied in [12] and [13], and shown under suitable conditions to converge to the Brownian web when appropriately rescaled.

2 Proof of Theorem 1

This section is entirely devoted to the proof of Theorem 1. Let us fix a sequence $\delta = \delta_n = 1/\sqrt{n}$ of positive numbers going to zero as $n \rightarrow \infty$. We want to verify conditions I_1, B_1 and B_2 of Theorem 2 for $\mathcal{X}_{\delta_n}$. Due to the translation invariance of the model, B_1 follows from I_1 (as will be argued below). So we only have to verify B_2 and I_1 . This will be done in Subsections 2.1 and 2.2 respectively. As a tool for both this verifications, we start by deriving a bound on the time of coalescence of two paths of the drainage model.

Coalescing random paths Let $u, v \in \mathbb{Z}^2$ be such that $u(1) \leq v(1)$ and $u(2) = v(2)$. Consider X^u and X^v and, for $t \geq u(2)$, define

$$Z_t = Z_t(u, v) = X_t^v - X_t^u, \quad (2.1)$$

Notice that $Z_{u(2)} = v(1) - u(1)$.

Remark 3 *As argued in [1] (see 3.1 in the proof of Theorem 2.1), $\{Z_t(u, v) : t \geq u(2)\}$ is a nonnegative martingale in $\mathbb{L}^2$. Also, $Z_t(u, v) \rightarrow 0$ a.s. as $t \rightarrow \infty$.*

Estimates for the tail of coalescence times We need to control the tail of the meeting time of two coalescing random paths starting at the same time

at a distance 1 apart. Define

$$\tau := \min\{t \geq 0 : Z_t = 0\} \quad (2.2)$$

Theorem 4 *There exists a constant $c_2 > 0$, such that $\mathbb{P}(\tau > t) \leq c_2/\sqrt{t}$.*

In order to prove Theorem 4 it is enough to consider the case in which $u = (0, 0)$ and $v = (1, 0)$. By Remark 3, Skorohod representation holds: that it is to say, there exists a Brownian motion with coefficient of diffusion 1 starting from 1 and stopping times $0 = T_0, T_1, T_2, \dots$ satisfying:

$$Z_t \stackrel{d}{=} B(T_t). \quad (2.3)$$

where $0 = T_0, T_1, T_2, \dots$ are such that

$$T_t = \inf\{s \geq T_{t-1} : B(s) - B(T_{t-1}) \notin (U_t(B(T_{t-1})), V_t(B(T_{t-1})))\}, \quad (2.4)$$

where $\{(U_t(m), V_t(m))\}, t \geq 1, m \in \mathbb{Z}\}$ is a family of random independent vectors and for all $m \in \mathbb{Z}$, $(U_t(m), V_t(m)) \stackrel{d}{=} (U_{t'}(m), V_{t'}(m))$ and $(U_t(m), V_t(m)) \in \{(0, 0), A\}$ with $A = \{\dots, -2, -1\} \times \{1, 2, \dots\}$. Note that since $\mathbb{P}(Z_t \geq 0) = 1$ we have that

$$\mathbb{P}(U_1(m) \geq -m) = 1, \forall m. \quad (2.5)$$

The following result will be needed later on.

Lemma 5 *For every $p < 1$, there exists a constant $c_1 \in (0, 1)$, such that for $m \geq 1$*

$$\mathbb{P}((U_1(m), V_1(m)) = (0, 0)) \leq c_1. \quad (2.6)$$

Proof

From the Skorohod representation, one readily concludes that the left hand side of (2.6) can be written as $\mathbb{P}(X_1^{(m,0)} - X_1^{(0,0)} = m)$. By conditioning on $X^{(0,0)}$, a straightforward computation yields

$$p^2 + \frac{1 - q^2}{2(1 + q^2)} q^2 \quad (2.7)$$

as an upper bound for that probability for all m , where $q = 1 - p$, and this expression is strictly less than 1 for $p < 1$. $\square$

Proof of Theorem 4

Let $\tau' := \min\{t \geq 0 : B(t) = 0\}$. From equations (2.3) and (2.5) we get

$$\mathbb{P}(\tau > t) = \mathbb{P}(\tau' > T_t). \quad (2.8)$$

Now, for $\zeta > 0$

$$\mathbb{P}(\tau' > T_t) \leq \mathbb{P}(\tau' > \zeta t) + \mathbb{P}(T_t \leq \zeta t) \leq \frac{c}{\sqrt{t}} + \mathbb{P}(T_t \leq \zeta t), \quad (2.9)$$

where $c = c(\zeta) \in (0, \infty)$, and the second inequality follows by the well known behavior of the tail of the distribution of τ' . By the Markov inequality, we have

$$\mathbb{P}(T_t \leq \zeta t) = \mathbb{P}(e^{-\lambda T_t} \geq e^{-\lambda \zeta t}) \leq e^{\lambda \zeta t} \mathbb{E}(e^{-\lambda T_t}). \quad (2.10)$$

We now write

$$T_t = \sum_{i=1}^t S_i(Z_{i-1}),$$

where $(S_i(k), i \geq 1, k \in \mathbb{Z})$ are independent random variables. (Note that for $k \in \mathbb{Z}$, $(S_i(k), i \geq 1)$ are not identically distributed.) Then, for $\lambda > 0$

$$\begin{aligned} \mathbb{E}(e^{-\lambda T_t}) &= \mathbb{E} \left[\mathbb{E} \left(\exp \left\{ -\lambda \sum_{i=1}^{t-1} S_i(Z_{i-1}) \right\} \exp \{ -\lambda S_t(Z_{t-1}) \} \middle| \mathcal{F}_{t-1} \right) \right] \\ &\leq \mathbb{E} \left(\exp \left\{ -\lambda \sum_{i=1}^{t-1} S_i(Z_{i-1}) \right\} \right) \sup_{m \in \mathbb{Z}} \mathbb{E}(e^{-\lambda S(m)}) \\ &\leq \left[\sup_{m \in \mathbb{Z}} \mathbb{E}(e^{-\lambda S(m)}) \right]^t, \end{aligned} \quad (2.11)$$

where $\mathcal{F}_t$ is the σ -algebra generated by $\{Z_0, Z_1, \dots, Z_t\}$.

Hence,

$$\mathbb{P}(T_t \leq \zeta t) \leq \left[e^{\lambda \zeta} \sup_{m \in \mathbb{Z}} \mathbb{E}(e^{-\lambda S(m)}) \right]^t, \quad (2.12)$$

and by Lemma 5 and Skorohod representation, we get (ommiting m 's)

$$\begin{aligned} \mathbb{E}(e^{-\lambda S}) &\leq \mathbb{P}((U, V) = (0, 0)) \\ &+ \mathbb{E}(e^{-\lambda S} | (U, V) = (-1, 1)) (1 - \mathbb{P}((U, V) = (0, 0))) \\ &= \left(1 - \frac{1}{\cosh(\sqrt{2\lambda})} \right) \mathbb{P}((U, V) = (0, 0)) + \frac{1}{\cosh(\sqrt{2\lambda})} \\ &\leq \left(1 - \frac{1}{\cosh(\sqrt{2\lambda})} \right) c_1 + \frac{1}{\cosh(\sqrt{2\lambda})}. \end{aligned} \quad (2.13)$$

Indeed, we first notice that S is the hitting time of (U, V) by B , and thus the case $(U, V) = (-1, 1)$ is dominated by the cases where $(U, V) \neq (0, 0)$; this justifies the first inequality in (2.13). The equality is a well known result (see e.g. Theorem 5.7 in [11]), and the last inequality follows from Lemma 5. Since $c_1 < 1$ (uniformly in m), we may find $\lambda_0 > 0$ and $\zeta_0 > 0$ such that

$$c_3 := \left(1 - \frac{1}{\cosh(\sqrt{2\lambda_0})} \right) c_1 + \frac{1}{\cosh(\sqrt{2\lambda_0})} < e^{-\lambda_0 \zeta_0}. \quad (2.14)$$

Then, from (2.12)

$$\mathbb{P}(T_t < \zeta_0 t) \leq (c_3 e^{\lambda_0 \zeta_0})^t := c_4^t, \quad (2.15)$$

where, from (2.14), c_4 may be taken < 1 . It follows that there exists $c_5 \in (0, \infty)$ such that $c_4^t \leq c_5/\sqrt{t}$. Making now $c_2 = c + c_5$, the result follows from (2.9) and (2.15). $\square$

2.1 Verification of condition B₂

The drainage network model is translation invariant, then we can eliminate $\sup_{(a,t_0) \in \mathbb{Z}^2}$ in (B₂) and consider $a = t_0 = 0$.

Then, to prove (B₂) it suffices to show that

$$\epsilon^{-1} \limsup_{N \rightarrow \infty} \mathbb{P} \left(\eta_{\mathcal{X}}(0, tN, 0, \epsilon\sqrt{N}) \geq 3 \right) \rightarrow 0, \text{ if } \epsilon \rightarrow 0+. \quad (2.16)$$

For $j \in \mathbb{Z} \cap [0, \epsilon\sqrt{N}]$ let $X_j = \{X^{(j,0)}(k), k \in \mathbb{Z} \cap [0, tN]\}$ be the $\lfloor tN \rfloor$ -step trajectory starting in $(j, 0)$.

We now introduce the counting variable $\eta' = |\{X_j(tN) : 1 \leq j \leq n\}|$, where $X_j(tN) = X^{(j,0)}(\lfloor tN \rfloor)$. Then,

$$\mathbb{P} \left(\eta_{\mathcal{X}}(0, tN; 0, \epsilon\sqrt{N}) \geq 3 \right) = \mathbb{P}(\eta' \geq 3). \quad (2.17)$$

Since

$$\{\eta' \geq 3\} = \cup_{j=1}^{n-1} \{X_{j-1}(tN) < X_j(tN) < X_n(tN)\},$$

where n is short for $\lfloor \epsilon\sqrt{n} \rfloor$, we have that

$$\begin{aligned} \mathbb{P}(\eta' \geq 3) &\leq \sum_{j=1}^{n-1} \mathbb{P}(X_{j-1}(tN) < X_j(tN) < X_n(tN)) \\ &= \sum_{j=1}^{n-1} \int_{\bar{\Pi}_j} \mathbb{P}(X_{j-1}(tN) < X_j(tN) < X_n(tN) | X_j = \pi) \mathbb{P}(X_j = \pi) \\ &= \sum_{j=1}^{n-1} \int_{\bar{\Pi}_j} \mathbb{P}(X_{j-1}(tN) < X_j(tN) | X_j = \pi) \\ &\quad \cdot \mathbb{P}(X_j(tN) < X_n(tN) | X_j = \pi) \mathbb{P}(X_j = \pi), \end{aligned} \quad (2.18)$$

where $\bar{\Pi}_j$ stands for the state space of X_j . The last equality follows from the fact that given $X_j = \pi$, the events $\{X_{j-1}(tN) < X_j(tN)\}$ and $\{X_j(tN) < X_n(tN)\}$ are independent.

At this point we want to appeal to the Harris-FKG inequality. So we must establish monotonicity properties of the conditional probabilities on the right of (2.18).

We begin by introducing a partial order $\prec$ on $\bar{\Pi}_j$ as follows. Given π_1 and $\pi_2 \in \bar{\Pi}_j$, say that

$$\pi_1 \prec \pi_2 \Leftrightarrow \pi_1(\ell) - \pi_1(k) \leq \pi_2(\ell) - \pi_2(k), \quad (2.19)$$

for every $\ell \geq k \geq 0, \ell, k \in \mathbb{Z} \cap [0, tN]$.

The partial order we have just introduced is an order on the increments of the trajectories belonging to events in $\bar{\Pi}_j$. Then, if the increments are increasing, the corresponding events will also be increasing.

Proposition 6 *Let $\pi_1, \pi_2 \in \bar{\Pi}_j$ be such that $\pi_1 \prec \pi_2$. Then, for $k < j < n$*

$$\mathbb{P}[X_k(tN) < X_j(tN) | X_j = \pi_1] \leq \mathbb{P}[X_k(tN) < X_j(tN) | X_j = \pi_2], \quad (2.20)$$

$$\mathbb{P}[X_j(tN) < X_n(tN) | X_j = \pi_1] \geq \mathbb{P}[X_j(tN) < X_n(tN) | X_j = \pi_2]. \quad (2.21)$$

Now, using Proposition 6, and since the increments of X_j are independent, we may apply Harris-FKG and we find an upper bound for (2.18) as follows.

$$\begin{aligned}
& \sum_{j=1}^{n-1} \int_{\bar{\Pi}_j} \mathbb{P}(X_{j-1}(tN) < X_j(tN) | X_j = \pi) \mathbb{P}(X_j = \pi) \\
& \quad \cdot \int_{\bar{\Pi}_j} \mathbb{P}(X_j(tN) < X_n(tN) | X_j = \pi) \mathbb{P}(X_j = \pi) \\
& = \sum_{j=1}^{n-1} \mathbb{P}(X_{j-1}(tN) < X_j(tN)) \mathbb{P}(X_j(tN) < X_n(tN)) \\
& \leq \sum_{j=1}^{n-1} \mathbb{P}(X_{j-1}(tN) < X_j(tN)) \mathbb{P}(X_0(tN) < X_n(tN)) \\
& \leq n \mathbb{P}(X_0(tN) < X_1(tN)) \mathbb{P}(X_0(tN) < X_{\epsilon\sqrt{N}}(tN)) \\
& \leq \epsilon\sqrt{N} \mathbb{P}(\tau > tN) \mathbb{P}(\tau_{\epsilon,N} > tN),
\end{aligned}$$

where τ is the coalescing time between the trajectories of $X^{(0,0)}$ and $X^{(1,0)}$ and $\tau_{\epsilon,N}$ is the coalescing time for $X^{(0,0)}$ and $X^{(n,0)}$.

From (2.17) we get

$$\epsilon^{-1} \mathbb{P}(\eta_\chi(0, tN; 0, \epsilon\sqrt{N}) \geq 3) \leq \sqrt{N} \mathbb{P}(\tau > tN) \mathbb{P}(\tau_{\epsilon,N} > tN). \quad (2.22)$$

From Theorem 8, we have that

$$\limsup_{N \rightarrow \infty} \mathbb{P}(\tau_{\epsilon,N} > tN) = \mathbb{P}(\tau_{\epsilon,B} > t),$$

where $\tau_{\epsilon,B}$ is the coalescing time between two Brownian motions starting at a distance ϵ at time zero. Also, recall that $\mathbb{P}(\tau_{\epsilon,B} > t)$ is well known to be $O(\epsilon)$. On the other hand, by Lemma 4, there exists a constant $c'_2 > 0$ such that $\mathbb{P}(\tau > tN) < c'_2/\sqrt{tN}$. Taking $\limsup_{N \rightarrow \infty}$ in both sides of (2.22) we get

$$\epsilon^{-1} \limsup_{n \rightarrow \infty} \mathbb{P}(\eta_\chi(0, tN; 0, \epsilon\sqrt{N}) \geq 3) = O(\epsilon),$$

which shows (2.16).

Proof of Proposition 6

It is enough to argue (2.20), since by the symmetry of the model we have

$$\mathbb{P}[X_j(tN) < X_n(tN) | X_j = \pi] = \mathbb{P}[X_{2j-n}(tN) < X_j(tN) | X_j = \pi^-], \quad (2.23)$$

where π^- is the path of $\bar{\Pi}_j$ whose increments are the opposite of the respective ones of π .

The next reduction is that it is enough to consider π_1, π_2 such that $\pi_1(s) = \pi_2(s)$ for $0 \leq s < t_0$ and $\pi_2(s) = \pi_1(s) + 1$ for $s \geq t_0$ for some $t_0 \in [0, tN] \cap \mathbb{Z}$. From now on, we will be dealing with such pair of paths π_1 and π_2 .

Remark 7 Given that $X_j = \pi$, the distribution of (Ω, Υ) changes as follows. Let $I_0 = (j, 0)$ and for $k \in [1, tN] \cap \mathbb{Z}$, let I_k be the interval of integer numbers with end points $2\pi(k-1) - \pi(k)$ and $\pi(k)$. (I_k can be a single point.) Then, off $\cup_{k=0}^{\lfloor tN \rfloor} I_k$, the distribution of (Ω, Υ) does not change. In $\cup_{k=0}^{\lfloor tN \rfloor} I_k$, the variables of (Ω, Υ) are independent of the ones off $\cup_{k=0}^{\lfloor tN \rfloor} I_k$, and they are also independent among themselves, except the pairs $(\omega(2\pi(k-1) - \pi(k), k), v(\pi(k-1), k-1))$ for which $|I_k| > 1$, whose distribution is described below. First, for each $k \in [1, tN] \cap \mathbb{Z}$, $\omega(\pi(k), k) = 1$. If $|I_k| > 1$, then $\omega(z_1, k) = 0$ for z_1 in I_k minus its endpoints, and $\omega(2\pi(k-1) - \pi(k), k)$ is a Bernoulli random variable with parameter $p' = p/(2-p)$; given that $\omega(2\pi(k-1) - \pi(k), k) = 0$, then $v(\pi(k-1), k-1)$ is Bernoulli with parameter $1/2$; otherwise it is 0 or 1 depending on whether $\pi(k) - \pi(k-1) > 0$ or < 0 .

A further point is that given that $X_j = \pi$, the event $\{X_k(tN) < X_j(tN)\}$ depends on Ω only to the left of and including π , and on Υ only strictly to the left of π .

In order to prove (2.20), we couple the distributions of (Ω, Υ) given π_1 and π_2 conveniently (in the relevant region of $\mathbb{Z}^2$). Let us denote by $(\omega_i(z), v_i(z))$, $z \in \mathbb{Z}^2$, the general term of (Ω, Υ) given π_i , $i = 1, 2$, respectively. Then $(\omega_2(z), v_2(z)) = (\omega_1(z), v_1(z))$ for $0 \leq z_2 < t_0$, and $(\omega_2(z), v_2(z)) = (\omega_1(z_1 - 1, z_2), v_1(z_1 - 1, z_2))$ for $0 \leq z_2 > t_0$. The coupling on $z_2 = t_0$ will be different in different cases.

Case 1: $\pi_1(t_0) < \pi_1(t_0 - 1)$.

In this case, we make $(\omega_2(z_1, t_0), v_2(z_1, t_0)) = (\omega_1(z_1 - 1, t_0), v_1(z_1 - 1, t_0))$ for $z_1 \leq \pi_2(t_0)$, and $(\omega_2(z_1, t_0), v_2(z_1, t_0)) = (\omega_1(z_1 + 1, t_0), v_1(z_1 + 1, t_0))$ for $z_1 > \pi_2(t_0)$.

One readily checks in this case that for every realization $(\omega_1, v_1)(\cdot)$ of (Ω, Υ) given $X_j = \pi_1$, $(\omega_2, v_2)(\cdot)$ given by the above coupling is a realization of (Ω, Υ) given $X_j = \pi_2$, and that

$$\{(\omega_1, v_1)(\cdot) : X_k(tN) < X_j(tN)\} \subset \{(\omega_2, v_2)(\cdot) : X_k(tN) < X_j(tN)\}. \quad (2.24)$$

This then establishes (2.20) in this case.

Case 2: $\pi_1(t_0) \geq \pi_1(t_0 - 1)$.

We start with a coupling which is the symmetric of the one above. It does not quite give us (2.24), but something nevertheless useful.

Auxiliary coupling:

Given $(\omega_2, v_2)(\cdot)$, we make $(\omega_1(z_1, t_0), v_1(z_1, t_0)) = (\omega_2(z_1 + 1, t_0), v_2(z_1 + 1, t_0))$ for $z_1 \geq \pi_1(t_0)$, and $(\omega_1(z_1, t_0), v_1(z_1, t_0)) = (\omega_2(z_1 - 1, t_0), v_2(z_1 - 1, t_0))$ for $z_1 < \pi_1(t_0)$. One can readily check that this provides a coupling of (Ω, Υ) given $X_j = \pi_1$ and (Ω, Υ) given $X_j = \pi_2$. It is possible to find realizations of $(\omega_2, v_2)(\cdot)$ for which with this coupling (2.20) does not hold. But the following can be readily checked.

$$\{(\omega_1, v_1)(\cdot) : X_k(t_0) < X_j(t_0)\} \subset \{(\omega_2, v_2)(\cdot) : X_k(t_0) < X_j(t_0)\}. \quad (2.25)$$

This will be used in the coupling discussed next.

We begin by pointing out that we could not indeed find a direct coupling for which (2.24) holds, so we take an indirect route. We start by considering $D_0 = X_j(t_0) - X_k(t_0)$. In order to insure the existence of a coupling of $(\omega_1, v_1)(\cdot)$

and $(\omega_2, v_2)(\cdot)$ satisfying (2.24), it is enough to show that the distribution of D_0 given $X_j = \pi_2$ dominates that of D_0 given $X_j = \pi_1$. Indeed, if this holds, then we can find a coupling of the two distributions such that the domination takes place almost surely. The respective random variables depend of course only on the history up to t_0 , and thus we are free to choose any coupling of $(\omega_1, v_1)(\cdot)$ and $(\omega_2, v_2)(\cdot)$ above t_0 , and choosing as in Case 1, we readily get (2.24).

In order to establish the above mentioned domination, we first define $\Delta = 2(\pi_1(t_0) - \pi_1(t_0 - 1))$ and $\tilde{D}_0 = \pi_1(t_0) - \Delta - X_k(t_0)$. Denote by D_i , $i = 1, 2$, the random variable whose distribution equals that of the conditional distribution of $\tilde{D}_0$ given that $X_j = \pi_i$, $i = 1, 2$, respectively. Then the above mentioned domination is equivalent to

$$\mathbb{P}(D_2 \geq k - 1) \geq \mathbb{P}(D_1 \geq k), \quad k \geq 0. \quad (2.26)$$

We will argue (2.26) using different couplings for different cases. The case $k = 0$ of (2.26) is established as follows. If $\pi_1(t_0) = \pi_1(t_0 - 1)$, there is nothing to prove. If $\pi_1(t_0) > \pi_1(t_0 - 1)$, then $D_1 \geq 0$ is equivalent to the left hand side of (2.25), and $D_2 \geq -1$ is equivalent to $D_2 \geq 0$, which is in turn equivalent to the right hand side of (2.25). So the auxiliary coupling can be used in this case. It can also be used if $k = 1$. Indeed, in this case, $D_2 \geq 0$ is equivalent to the right hand side of (2.25) and $D_1 \geq 1$ is either equivalent to the left hand side of (2.25) (if $\pi_1(t_0) = \pi_1(t_0 - 1)$), or is in any case contained in it.

For $k \geq 2$, we will resort to another coupling, described as follows.

1. For $z_1 > \pi_1(t_0)$, $(\omega_2(z_1, t_0), v_2(z_1, t_0)) = (\omega_1(z_1 - 1, t_0), v_1(z_1 - 1, t_0))$;
2. for $\pi_1(t_0) - 2(\pi_1(t_0) - \pi_1(t_0 - 1)) \leq z_1 \leq \pi_1(t_0)$, $\omega_2(z_1, t_0) = 0$ and $v_2(z_1, t_0)$ are i.i.d. Bernoullis with parameter $1/2$ independent of all else;
3. for $z_1 = \pi_1(t_0) - 2(\pi_1(t_0) - \pi_1(t_0 - 1)) - 1$, $(\omega_2(z_1, t_0), v_2(z_1, t_0)) = (\xi\omega_1(z_1, t_0), \tilde{v})$, where ξ is a Bernoulli with parameter $1/(2 - p)$ independent of all else, and $\tilde{v}$ depends only on $\xi\omega_1(z_1, t_0)$; if the latter random variable vanishes, then $\tilde{v}$ is Bernoulli with parameter $1/2$ independent of all else; otherwise, $\tilde{v} = 1$;
4. and for $z_1 < \pi_1(t_0) - 2(\pi_1(t_0) - \pi_1(t_0 - 1)) - 1$, $(\omega_2(z_1, t_0), v_2(z_1, t_0)) = (\omega_1(z_1, t_0), v_1(z_1, t_0))$.

One readily checks that for every realization $(\omega_1, v_1)(\cdot)$ of (Ω, Υ) given $X_j = \pi_1$, $(\omega_2, v_2)(\cdot)$ given by the above coupling is a realization of (Ω, Υ) given $X_j = \pi_2$, and that

$$\{(\omega_1, v_1)(\cdot) : D_1 \geq k\} \subset \{(\omega_2, v_2)(\cdot) : D_2 \geq k\} \quad (2.27)$$

in the remainder cases. $\square$

2.2 Weak convergence to coalescing Brownian motions

The main purpose of this part of the paper is to prove condition *I1* of Theorem 2. Also, we prove condition *B1*, which is in fact a consequence of condition *I1* for this model.

Denote a single trajectory belonging to the drainage network model and starting from $a \in \mathbb{Z}$ at time $j \in \mathbb{Z}$ by $X^{a,j} = \{X^{a,j}(l), l \geq j\}$. Then,

Theorem 8 Let $(y_0, s_0), (y_1, s_1), \dots, (y_k, s_k)$ be $k+1$ different points in $\mathbb{R}^2$ such that $s_0 \leq s_1 \leq \dots \leq s_k$ and if $s_{i-1} = s_i$ for some $i, i = 1, \dots, k$, then $y_{i-1} < y_i$. If $Z_n^{(i)} = \{Z_n^{(i)}(t) := n^{-1/2} X_n^{(i)}(\lfloor nt \rfloor) := n^{-1/2} X^{\lfloor y_i \sqrt{n} \rfloor, \lfloor s_i n \rfloor}(\lfloor nt \rfloor), t \geq s_i\}$; then

$$\{Z_n^{(i)}, i = 0, \dots, k\} \Rightarrow \{W^{(i)}, i = 0, \dots, k\}, \quad (2.28)$$

as $n \rightarrow \infty$, where the $W^{(i)}$ are $k+1$ coalescing Brownian motions with positive diffusion coefficient σ , starting at $\{(y_0, s_0), \dots, (y_k, s_k)\}$.

Remark 9 Here, $\Rightarrow$ stands for weak convergence in Π^{k+1} (the product of $k+1$ copies of Π , the path space (see [5])).

Remark 10 σ^2 is the variance of $X^{0,0}(1)$, which one readily computes from its distribution, which is in turn straightforward to obtain, as $\frac{q(1+q^2)}{p^2(1+q)^2}$.

Proof of Theorem 8

The proof is divided in three parts. The arguments are somewhat standard, so we will be sketchy at a few points.

Part I

When $k = 0$, the result follows from Donsker' invariance principle for single paths since $X^{a,j}$ is a random walk with variance σ^2 .

Part II

In this part, we consider the case where $k \geq 1$ and $s_0 = \dots = s_k$ (and $y_{i-1} < y_i$ for every $i, i = 1, \dots, k$). The general case will be treated on Part III. Since the increments are stationary and spatially homogeneous we will take $s_0 = y_0 = 0$.

a) We first treat the case $k = 1$. Without loss of generality, we suppose that we are working in a probability space in which

$$Z_n^{(1)} \xrightarrow[n \rightarrow \infty]{a.s.} W^{(1)}. \quad (2.29)$$

The following proposition, besides implying the result in the case $k = 1$, is also the building block of the argument for the other cases of this part.

Proposition 11 The conditional distribution of $Z_n^{(0)}$ given $Z_n^{(1)}$ converges almost surely to the distribution of $W^{(0)}$ given $W^{(1)}$.

Proof

The strategy is to approximate the paths $Z_n^{(0)}$ and $Z_n^{(1)}$ before the time when they first come close (at a distance of order n^α , $\alpha < 1/2$) by independent paths; then to show that after that time they meet quickly. Let us write

$$X_n^{(i)}(l) = \lfloor y_i \sqrt{n} \rfloor + \sum_{h=1}^l S_h^{(i,n)}, \quad (2.30)$$

where for each $i = 0, 1$ and $n \geq 1$, $S_h^{(i,n)}, h \geq 1$, are i.i.d. random variables distributed as $X^{0,0}(1)$. Next, we introduce independent copies of $S_h^{(i,n)}, i = 0, 1, n \geq 1$. For each $n \geq 1$, let $\{\tilde{S}_h^{(i,n)}, h \geq 1, i = 0, 1\}$ be i.i.d. random variables such that $\tilde{S}_1^{(0,n)} \sim X^{0,0}(1)$ and define random variables $\hat{S}_h^{(i,n)}$ by

$$\hat{S}_h^{(i,n)} = \begin{cases} S_h^{(i,n)}, & \text{if } |S_h^{(i,n)}| \leq n^\alpha \\ \tilde{S}_h^{(i,n)}, & \text{otherwise,} \end{cases} \quad (2.31)$$

where α is a positive number to be specified later. Moreover, for $l \geq 1$, let $\hat{X}_n^{(i)}(l) = \lfloor y_i \sqrt{n} \rfloor + \sum_{h=1}^l \hat{S}_h^{(i,n)}$. Define

$$\hat{\tau}_n = \min\{l \geq 1 : \hat{X}_n^{(1)}(l) - \hat{X}_n^{(0)}(l) \leq 3n^\alpha\} \quad (2.32)$$

and

$$\tau_n = \min\{l \geq 1 : X_n^{(1)}(l) - X_n^{(0)}(l) \leq 3n^\alpha\}, \quad (2.33)$$

and let

$$\tilde{X}_n^{(i)}(l) = \begin{cases} \hat{X}_n^{(i)}(l), & \text{if } l \leq \hat{\tau}_n \\ \hat{X}_n^{(i)}(\hat{\tau}_n) + \sum_{h > \hat{\tau}_n} \check{S}_h^{(i,n)}, & \text{otherwise,} \end{cases} \quad (2.34)$$

where

$$\check{S}_h^{(i,n)} = \begin{cases} \tilde{S}_h^{(0,n)}, & \text{if } i = 0 \\ S_h^{(1,n)}, & \text{if } i = 1. \end{cases} \quad (2.35)$$

From the exponential tail of the distribution of $S_h^{(i,n)}$ (which is independent of i) we readily get the following result.

Lemma 12 *Given $\beta < \infty$, the complement of the event*

$$A_n := \{S_h^{(i,n)} = \hat{S}_h^{(i,n)}, i = 1, \dots, n^\beta, i = 0, 1\},$$

has probability going to zero superpolynomially fast as $n \rightarrow \infty$.

Corollary 13 *The event $B_n := \{X_n^{(i)}(l) = \hat{X}_n^{(i)}(l); l = 1, \dots, n^\beta; i = 0, 1\}$, is such that $\mathbb{P}[B_n^c] \rightarrow 0$ converges to superpolynomially fast as $n \rightarrow \infty$.*

From the construction, the properties stated in the following result are readily verified.

Lemma 14 $\tilde{X}_n^{(0)} := \{\tilde{X}_n^{(0)}(l), l \geq 1\}$ and $\tilde{X}_n^{(1)} := \{\tilde{X}_n^{(1)}(l), l \geq 1\}$ are independent processes and $X_n^{(i)} \sim \tilde{X}_n^{(i)}$ for each fixed $i = 0, 1$.

Corollary 15 *Let $\tilde{Z}_n^{(i)}(t) := n^{-1/2} \tilde{X}_n^{(i)}(\lfloor nt \rfloor)$, $t \geq 0$. Then,*

1. $\tilde{Z}_n^{(1)} \rightarrow W^{(1)}$ almost surely as $n \rightarrow \infty$;
2. $(\tilde{Z}_n^{(0)}, \tilde{Z}_n^{(1)}) \Rightarrow (\tilde{W}^{(0)}, W^{(1)})$, where $\tilde{W}^{(0)} \sim W^{(0)}$, and $\tilde{W}^{(0)}$ and $W^{(1)}$ are independent;
3. the conditional distribution of $\{\tilde{Z}_n^{(0)}(t), t \leq \frac{\hat{\tau}_n}{n}\}$ given $\tilde{Z}_n^{(1)}$ converges almost surely to that of $\{\tilde{W}^{(0)}(t), t \leq \tau\}$ given $W^{(1)}$, where $\tau = \inf\{t \geq 0 : \tilde{W}^{(0)}(t) = W^{(1)}(t)\}$.

Corollary 16 *The conditional distribution of $\{Z_n^{(0)}(l), l \leq \tau_n\}$ given $Z_n^{(1)}$ converges almost surely to that of $\{\tilde{W}^{(0)}(t), t \leq \tau\}$ given $W^{(1)}$.*

Lemma 17 *For $\alpha < 1/2$, there exists $\gamma = \gamma(\alpha) < 1$ such that, as $n \rightarrow \infty$,*

$$\mathbb{P}[X_n^{(0)}(l) \neq X_n^{(1)}(l), l = \tau_n, \dots, \tau_n + n^\gamma] \rightarrow 0. \quad (2.36)$$

Corollary 18 $\{Z_n^{(0)}(t), t \geq \frac{\tau_n}{n}\}$ converges in probability to $\{W^{(1)}(t), t \geq \tau\}$.

Then, Proposition 11 follows directly from Corollaries 16 and 18, and the Markov property. $\square$

Proof of Corollary 15

1. Let $\beta > 7/2$ be fixed. We claim that there exists a finite constant C and $\theta > 1$ such that

$$\mathbb{P}[\hat{\tau}_n > n^\beta] \leq C/n^\theta. \quad (2.37)$$

To argue this we first notice that $Y^{(n)} := \hat{X}_n^{(1)} - \hat{X}_n^{(0)}$ is a random walk on the integers starting from $\lfloor y_i \sqrt{n} \rfloor$ and with increments distributed as $Y^{(n)}(1) = \hat{S}_1^{(1,n)} - \hat{S}_1^{(0,n)}$. One may readily notice then that $\hat{\tau}_n$ is dominated by the sum of $\lfloor y_i \sqrt{n} \rfloor$ copies of $\tilde{\tau}_n$, the hitting time of the negative integers by a random walk with increments distributed as $Y^{(n)}(1)$, started at the origin. We then have that

$$\mathbb{P}(\hat{\tau}_n > n^\beta) \leq c_1 \sqrt{n} \mathbb{P}(\tilde{\tau}_n > c_2 n^{\beta-1/2}) \leq C/n^{(\beta-3/2)/2}, \quad (2.38)$$

where the first bound is straightforward from the above domination, and the last one follows from a standard fact on the tail of the distribution of hitting times for one dimensional random walks with finite second moment, like $Y^{(n)}$. The claim is thus justified.

We can then use (2.29), Corollary 13, and (2.37), together with Borel-Cantelli, to conclude.

2. Immediate from Lemma 17 and Donsker's theorem.
3. We take a version of $\tilde{Z}_n^{(0)}$, say $\bar{Z}_n^{(0)}$, such that

$$\left(\bar{Z}_n^{(0)}, \tilde{Z}_n^{(1)} \right) \xrightarrow[n \rightarrow \infty]{a.s.} \left(\bar{W}^{(0)}, W^{(1)} \right), \quad (2.39)$$

with $(\bar{W}^{(0)}, W^{(1)}) \sim (\tilde{W}^{(0)}, W^{(1)})$, and claim that $\bar{\tau}_n \rightarrow \bar{\tau}$ almost surely as $n \rightarrow \infty$, where $\bar{\tau}_n = \min\{t \geq 0 : \tilde{Z}_n^{(1)}(t) - \bar{Z}_n^{(0)}(t) \leq 3n^{\alpha-1/2}\}$ and $\bar{\tau} = \inf\{t \geq 0 : \bar{W}^{(0)}(t) = W^{(1)}(t)\}$. Indeed, the almost sure convergence of $\bar{\tau}_n$ follows from the property, that with probability one, two independent Brownian trajectories will instantaneously cross after touching, i.e.

$$\mathbb{P}[\forall \epsilon > 0, \bar{W}^{(0)}(t) > W^{(1)}(t) \text{ for some } t \leq \bar{\tau} + \epsilon] = 1,$$

and the following deterministic result (whose proof is an exercise). The convergence in distribution in the original space follows. $\square$

Lemma 19 *Let $f_n, g_n, f, g : \mathbb{R}^+ \rightarrow \mathbb{R}$ be continuous functions such that $f(0) < g(0)$, $T = \inf\{t \geq 0 : f(t) = g(t)\}$ is finite and has the property that for every $\delta > 0$ there exists $t \in [T, T + \delta]$ with $f(t) > g(t)$. Suppose in addition that $\lim_{n \rightarrow \infty} \sup_{0 \leq t \leq T+1} |f_n(t) - f(t)| = 0$, and $\lim_{n \rightarrow \infty} \sup_{0 \leq t \leq T+1} |g_n(t) - g(t)| = 0$, and let $T_n = \inf\{t \geq 0 : g_n(t) - f_n(t) \leq \epsilon_n\}$, where (ϵ_n) is a given sequence of numbers vanishing as $n \rightarrow \infty$. Then, $\lim_{n \rightarrow \infty} T_n = T$.*

Proof of Corollary 16

Immediate from Corollary 13 and 3. of Corollary 15. $\square$

Proof of Lemma 17

Let $\sigma_n = \inf\{l \geq \tau_n : X_n^{(0)}(l) = X_n^{(1)}(l)\} - \tau_n$. It can be readily checked that σ_n is dominated by $\tilde{\sigma}_n$, the hitting time of 0 by $Z_t((0, 0), ([3n^\alpha], 0))$ (see (2.1)). Proceeding as in the proof of Theorem 4, we find

$$\mathbb{P}(\tilde{\sigma}_n > n^\gamma) = \mathbb{P}(\sigma'_n > T_{n^\gamma}) \leq \mathbb{P}(\sigma'_n > \zeta n^\gamma) + \mathbb{P}(T_{n^\gamma} \leq \zeta n^\gamma), \quad (2.40)$$

where σ'_n is the hitting time of 0 by a Brownian motion started at $\lfloor 3n^\alpha \rfloor$ (see (2.8) and (2.9)). Choosing now γ in $(2\alpha, 1)$, one readily checks that the first term on the right of (2.40) vanishes as $n \rightarrow \infty$. That so does the second one follows by (2.15), and the proof is complete. $\square$

Proof of Corollary 18

By (2.29) and Lemma 17, it is enough to show that for every $\epsilon > 0$, as $n \rightarrow 0$

$$\mathbb{P}(\max_{0 \leq l \leq n^\gamma} (X_n^{(1)}(l + \tau_n) - X_n^{(0)}(l + \tau_n)) > \epsilon\sqrt{n}) \rightarrow 0. \quad (2.41)$$

By the Markov property and an elementary domination argument, the left hand side of (2.41) is bounded above by

$$\mathbb{P}(\max_{0 \leq l \leq n^\gamma} \hat{Z}_l^{(n)} > \epsilon\sqrt{n}), \quad (2.42)$$

where $\hat{Z}_l^{(n)} = Z_l((0, 0), (\lfloor 3n^\alpha \rfloor, 0))$ (see (2.1)). Now by Doob's inequality, (2.42) is bounded above by constant times $\mathbb{E}(\hat{Z}_{n^\gamma}^{(n)})/\sqrt{n} \leq 3n^{\alpha-1/2} \rightarrow 0$ as $n \rightarrow \infty$. $\square$

b) We now consider the case $k > 1$. It suffices to prove that

$$\mathbb{E}[f_0(Z_n^{(0)}) \dots f_k(Z_n^{(k)})] \xrightarrow{n \rightarrow \infty} \mathbb{E}[f_0(W^{(0)}) \dots f_k(W^{(k)})] \quad (2.43)$$

for any $f_0, \dots, f_k \in C_b(\Pi, \mathbb{R})$, the space of all real-valued and bounded functions defined on Π .

The left hand side of (2.43) is equivalent to

$$\mathbb{E}\{f_1(Z_n^{(1)})\mathbb{E}[f_0(Z_n^{(0)})f_2(Z_n^{(2)}) \dots f_k(Z_n^{(k)})|Z_n^{(1)}]\}. \quad (2.44)$$

It follows from Lemma 20 below that the conditional expectation equals

$$\mathbb{E}[f_0(Z_n^{(0)})|Z_n^{(1)}] \mathbb{E}[f_2(Z_n^{(2)}) \dots f_k(Z_n^{(k)})|Z_n^{(1)}].$$

Since $Z_n^{(1)} \xrightarrow{a.s.} W^{(1)}$ by the inductive hypothesis the conditional expectations above converge, almost surely, to $\mathbb{E}[f_0(W^{(0)})|W^{(1)}]$ and $\mathbb{E}[f_2(W^{(2)}) \dots f_k(W^{(k)})|W^{(1)}]$, respectively. On the other hand, by the dominated convergence theorem we get the convergence of (2.44) to

$$\mathbb{E}\{f_1(W^{(1)})\mathbb{E}[f_0(W^{(0)})|W^{(1)}] \mathbb{E}[f_2(W^{(2)}) \dots f_k(W^{(k)})|W^{(1)}]\},$$

which equals

$$\mathbb{E}\{f_1(W^{(1)})\mathbb{E}[f_0(W^{(0)})f_2(W^{(2)}) \dots f_k(W^{(k)})|W^{(1)}]\} \quad (2.45)$$

Since (2.45) equals the right hand side of (2.43), we have the result. $\square$

Lemma 20 For each $n \geq 1$, given $Z_n^{(1)}, Z_n^{(0)}$ is independent of $(Z_n^{(2)}, \dots, Z_n^{(k)})$.

Proof

$Z_n^{(0)}$ depends only on the variables of (Ω, Υ) to the left of and including $Z_n^{(1)}$, and $(Z_n^{(2)}, \dots, Z_n^{(k)})$ depends only on the variables of (Ω, Υ) to the right of and including $Z_n^{(1)}$. The result follows from the product structure of the conditional distribution of (Ω, Υ) given $Z_n^{(1)}$ (see Remark 7), and the facts that,

given $Z_n^{(1)}$, the variables of Ω along $Z_n^{(1)}$ are fixed, and when $Z_n^{(0)}$ depends on the variable of Υ in a particular position of $Z_n^{(1)}$ (that happens when, and only when, the increment of $Z_n^{(1)}$ from the given position is strictly positive), then $(Z_n^{(2)}, \dots, Z_n^{(k)})$ does not depend on that particular variable, and vice-versa. $\square$

Part III

We now take on the general case and proceed by induction on $k \geq 0$. The case $k = 0$ was argued in Part I above. Suppose now that for some $k \geq 1$, the result holds up to $k - 1$, and let $(y_0, s_0), \dots, (y_k, s_k)$ be as in the statement of Theorem 8. If the condition on (s_i) of Part II holds, then there is nothing to prove. Otherwise, there exists $i \in \{1, \dots, k\}$ such that $s_i > s_{i-1}$. Let i_0 be the maximum between such i 's. By the induction hypothesis,

$$(\{Z_n^{(i)}(t), s_i \leq t \leq s_{i_0}\}, 0 \leq i < i_0) \Rightarrow (\{W^{(i)}(t), s_i \leq t \leq s_{i_0}\}, 0 \leq i < i_0), \quad (2.46)$$

and we can suppose that such convergence occurs almost surely. By the Markov property and Part II (strengthened to accomodate the case where the y_i 's may depend on n and converge to, say, $\tilde{y}_i$ as $n \rightarrow \infty$; this requires minor changes in the argument above), we have that, given (2.46),

$$(\{Z_n^{(i)}(t), t \geq s_{i_0}\}, 0 \leq i \leq k+1) \Rightarrow (\{W^{(i)}(t), t \geq s_{i_0}\}, 0 \leq i \leq k+1). \quad (2.47)$$

Then, Theorem 8 follows from (2.46), (2.47) and the Markov property. $\square$

Verification of B_1

We note that $\eta_{\mathcal{X}_n}(0, t; 0, \epsilon) \geq 2$ if and only if the two trajectories starting at time 0 from the border of the interval $[0, \epsilon]$ have not met up to time t . By Theorem 8, the rescaled trajectories of a finite collection of paths belonging to the drainage network model converge to those of coalescing Brownian motion. Then, it is straightforward to get, for the drainage network model, that

$$\limsup_{n \rightarrow \infty} \mathbb{P}(\eta_{\mathcal{X}_n}(0, t; 0, \epsilon) \geq 2) = 2\Phi(\epsilon/\sqrt{2t}) - 1,$$

where $\Phi(\cdot)$ is the standard normal distribution function. This together with the translation invariance of the model concludes the proof.

Acknowledgements

This work started as part of the master's project of the third author and was completed during the post-doc project of the first one, both at IME-USP and financed by FAPESP. We thank Pablo Ferrari for discussions on the coupling argument of Subsection 2.1.

References

- [1] S. Gangopadhyay, R. Roy, A. Sarkar, Random oriented trees: a model of drainage networks, *Ann. App. Probab.* **14**, 1242-1266 (2004).
- [2] R. Arratia, Limiting point processes for rescalings of coalescing and annihilating random walks on $\mathbb{Z}^d$, *Ann. Prob.* **9**, pp 909-936 (1981).

- [3] R. Arratia, Coalescing Brownian motions and the voter model on $\mathbb{Z}$. Unpublished partial manuscript (circa 1981), available at rarratia@math.usc.edu.
- [4] B. Tóth, W. Werner, The true self-repelling motion, *Prob. Theory Related Fields* **111**, pp 375-452 (1998).
- [5] L. R. G. Fontes, M. Isopi, C. M. Newman, K. Ravishankar, The Brownian web: characterization and convergence, *Ann. Probab.* **32**, 2857-2883 (2004)
- [6] L. R. G. Fontes, M. Isopi, C. M. Newman, K. Ravishankar, The Brownian web, *Proc. Natl. Acad. Sci. USA* **99**, no. 25, 15888 – 15893 (2002).
- [7] P. A. Ferrari, C. Landim, H. Thorisson, Poisson trees, succession lines and coalescing random walks, *Ann. Inst. H. Poincar Probab. Statist.* **40**, 141-152 (2004).
- [8] P. A. Ferrari, L. R. G. Fontes, X.-Y. Wu, Two-dimensional Poisson trees converge to the Brownian web, *Ann. Inst. H. Poincar Probab. Statist.* **41**, 851–858 (2005)
- [9] A. E. Scheidegger, A stochastic model for drainage patterns into an intra-montane trench, *Bull. Ass. Sci. Hydrol.* **12**, 15-20 (1967).
- [10] I. Rodriguez-Iturbe, A. Rinaldo, *Fractal river basins: chance and self-organization*, Cambridge Univ. Press, New York (1997).
- [11] R. Durrett, *Probability: Theory and Examples*. Second Edition, Duxbury Press, 1996.
- [12] C. M. Newman, K. Ravishankar, R. Sun, Convergence of coalescing non-simple random walks to the Brownian web, *Electron. J. Probab.* **10**, 21–60 (2005)
- [13] S. Belhaouari, T. Mountford, R. Sun, G. Valle, Convergence results and sharp estimates for the voter model interfaces, *Electron. J. Probab.* **11**, 768–801 (2006)